



Smithsonian Tropical Research Institute
Instituto Smithsonian de Investigaciones Tropicales

FOR IMMEDIATE RELEASE

Contact: Dr. Jesús Mavárez, mavarezj@si.edu, from U.S. via satellite 202 786 2094 ext. 8842, or 011+507 212-8842
or Beth King, Science Interpreter, kingb@si.edu, 202 786 2094 ext 8216 or 011+507 212-8216

Butterfly speciation event

In a matter of months, butterflies sporting the yellow and red wing color pattern of a wild species were created through simple laboratory crosses of a wild species with a red wing pattern with a wild species with a yellow wing pattern, researchers report in the June 15, 2006 edition of the journal *Nature*.

“We recreated the evolutionary steps that may have given rise to *Heliconius heurippa*, a hybrid butterfly species, in the lab,” explains Jesús Mavárez, Molecular Evolution Fellow at the Smithsonian Tropical Research Institute.

Sexual encounters between species resulting in hybrid offspring may be common in nature. However, homoploid hybrid species—fully sexual hybrid species resulting from crosses between two different parent species—are still considered to be quite rare. In *Ragoletis* fruit flies, Swordtail fishes and African Cichlids there is growing evidence for homoploid hybrid speciation. This report of the laboratory creation of a hybrid butterfly nearly identical to a known wild species, and the observation that hybrid individuals don’t mate with either parent species, provides the most convincing case to-date for homoploid hybrid speciation in animals.

Co-author Mauricio Linares (University of the Andes, Colombia) had long suspected that *H. heurippa*, boasting red and yellow wing stripes, could be a hybrid species resulting from a cross between *H. cydno* and *H. melpomene*. He set up a long term research project to test this idea. Linares, with his students Camilo Salazar and Christian Salcedo, designed and performed a series of crosses between *H. cydno* and *H. melpomene*. The result was a homoploid sexual hybrid very similar to *H. heurippa*.

“We found that a wing pattern almost identical to that of the hybrid can be obtained in months-- just 3 generations of lab crosses between *H. cydno* and *H. melpomene*! Moreover, natural hybrids from San Cristóbal, Venezuela, show wing patterns very similar to *H. heurippa*, further supporting the idea of a hybrid origin for this species,” Mavárez explains.

Butterflies in the genus *Heliconius* (~60 species) exhibit a tremendous diversity of brightly coloured wing patterns. Strikingly beautiful red, orange, yellow and white markings on black wings clearly signal to birds and other predators that the butterflies are toxic.

SI-246-2006

Smithsonian Institution
Smithsonian Tropical Research Institute
Apartado 2072
Balboa, Ancón
República de Panamá
Tel. 507.212.8216
FAX 507.212.8148
Email: kingb@tivoli.si.edu



Smithsonian Tropical Research Institute Instituto Smithsonian de Investigaciones Tropicales

The wing pattern signaling system is so finely tuned that butterflies with even slight deviations in color pattern—confusing signals-- suffer from higher predation. Non-toxic butterfly species may copy common wing patterns from this group in order to take advantage of the predator's learned avoidance response (a process known as mimicry). Color patterns also serve as mating cues; these butterflies are extremely choosy about finding mates with their own, species or race-specific, wing pattern. For *Heliconius*, color is everything.

Heliconius heurippa males choose females with red and yellow stripes. They don't choose females of their parent species (*H. cydno* and *H. melpomene*) because these species have only a yellow or a red stripe, respectively. "If you cover the red or the yellow stripe of a bicoloured hybrid female, hybrid males no longer find her the least bit attractive," continues Mavárez, who documented the genetic relationships between the three species at the Smithsonian molecular biology laboratories in Panama.

The "weird" wing pattern of *H. heurippa* individuals makes them undesirable as mates for members of their parents' species, but attractive to each other—and that's almost all it takes for speciation to occur.

"We're seeing that mate choice may be a very strong force driving hybrid speciation," adds STRI Staff Scientist, Eldredge Bermingham.

The process of speciation usually occurs when one species splits into two, but there are some cases in which new species result from the mating between two good species. Wait a minute, you may be saying, isn't the definition of a species is that it can only produce viable offspring with members of its own species?

Yes, but. Usually when members of different species mate, their offspring die or are infertile. On occasion, polyploid hybrids—individuals with multiple copies of their parents' chromosomes, survive and establish a new lineage or species. Polyploid hybrid speciation often occurs in plants when polyploid individuals go on to reproduce asexually. Some lizards, frogs, fish and many invertebrates also pull this trick.

But hybrid matings may also produce normal, sexual offspring with the same number of chromosomes as their parents (homoploids). These offspring don't usually go on to become new species, because they backcross with individuals from their parent species.

At least two other species in the group, *H. timareta* and *H. pacheus*, are also proposed to result from *H. cydno*/*H. melpomene* hybridization. "We will feel very fortunate if there are more hybrid species in this group," Mavárez continues, "Cases of hybrid animal species are extremely rare and you're very lucky to have one in your preferred group. So, imagine our excitement about the possibility of finding several hybrid species in *Heliconius*! A whole new set of questions could be addressed." A new research project to identify the specific genetic

SI-246-2006

Smithsonian Institution
Smithsonian Tropical Research Institute
Apartado 2072
Balboa, Ancón
República de Panamá
Tel. 507.212.8216
FAX 507.212.8148
Email: kingb@tivoli.si.edu



Smithsonian Tropical Research Institute
Instituto Smithsonian de Investigaciones Tropicales

contribution of each parental species to the genome of *H. heurippa* promises to shed new light on this kind of hybrid speciation.

##

The Smithsonian Tropical Research Institute (STRI), a unit of the Smithsonian Institution, headquartered in Panama City, Panama, furthers our understanding of tropical nature and its importance to human welfare, trains students to conduct research in the tropics and promotes conservation by increasing public awareness of the beauty and importance of tropical ecosystems. www.stri.org

Ref: Mavarez, J., Salazar, C.A., Bermingham, E., Salcedo, C., Jiggins, C.D. , Linares, M. 2006 Speciation by hybridization in Heliconius butterflies. Nature, 15 June, 2006.

Photo captions and credits:

heurippa.jpg Heliconius heurippa, mounted specimen, hybrid species resulting from mating of *H. cydno* x *H. melpomene*, Credit: Christian Salcedo, University of Florida Gainesville

Heurippa_live_salcedo.jpg Heliconius heurippa, live specimen, hybrid species resulting from mating of *H. cydno* x *H. melpomene*, Credit: Christian Salcedo, University of Florida Gainesville

Hibrido cydnomelpomene.bmp Natural hybrid between *H. cydno* x *H. melpomene*, San Cristobal, Venezuela, Credit: Mauricio Linares, Universidad de Los Andes, Colombia

Jesus183b.jpg Jesus Mavarez, Credit: Marcos Guerra, Smithsonian Tropical Research Institute

SI-246-2006

Smithsonian Institution
Smithsonian Tropical Research Institute
Apartado 2072
Balboa, Ancón
República de Panamá
Tel. 507.212.8216
FAX 507.212.8148
Email: kingb@tivoli.si.edu