

A sediment record of trophic state change in an Arkansas (USA) reservoir

Byron Winston · Sonja Hausmann ·
Jaime Escobar · William F. Kenney

Received: 20 May 2013 / Accepted: 18 December 2013
© Springer Science+Business Media Dordrecht 2013

Abstract Reservoir sediments are used cautiously in paleolimnological studies because of dating uncertainties, possible sediment disturbances and even concerns that indicators of trophic status may behave differently in reservoirs as opposed to natural lakes. We measured loss on ignition (LOI), carbon to nitrogen ratio (C:N), diatom abundance, total nitrogen (TN), total phosphorus (TP), TN:TP ratio, and carbon and nitrogen isotopes ($\delta^{13}\text{C}$ and $\delta^{15}\text{N}$) in an 83-cm sediment core to track recent trophic status changes in Beaver Reservoir,

Northwest Arkansas, USA. Measurements showed that LOI, TN, TP and diatom abundance increased significantly from the bottom to the top of the core ($p < 0.001$). The C:N ratio and $\delta^{13}\text{C}$ indicated a predominantly algal source for organic matter in the sediments. Increases in TN and TP were positively correlated with human population growth ($p < 0.01$) and the TN:TP ratio recorded a shift from phosphorus to nitrogen limitation around 1990. This shift may have encouraged cyanobacterial growth that caused episodes of taste and odor problems in the reservoir. This study suggests that despite concerns about sediment dating and disturbance, reservoir sediments can provide valuable information on past water quality changes.

B. Winston (✉)
Department of Crop, Soil and Environmental Sciences,
University of Arkansas, 114 Plant Sciences Building,
Fayetteville, AR 72701, USA
e-mail: bwinsto@uark.edu

S. Hausmann
Academy of Natural Sciences of Drexel University,
1900 Benjamin Franklin Parkway, Philadelphia,
PA 19103, USA

J. Escobar
Departamento de Ingeniería Civil y Ambiental,
Universidad del Norte, Km 5 Via Puerto Colombia,
Barranquilla, Colombia

J. Escobar
Center for Tropical Paleocology and Archaeology,
Smithsonian Tropical Research Institute (STRI),
Panama City, Panama

W. F. Kenney
Land Use and Environmental Change Institute, University
of Florida, Gainesville, FL 32611, USA

Keywords Paleolimnology · Reservoir ·
Sediments · Water quality · Geochemistry

Introduction

Paleolimnological studies in the south-central United States are rare relative to the northern USA because the former region lacks natural, glacial lakes and because there are concerns that reservoir deposits may not accurately record water quality history (Shotbolt et al. 2005, 2006; Filstrup et al. 2010). Of particular concern are the possibilities of sediment disturbance and the relatively young ages of reservoirs, which may hinder application of traditional dating techniques

(Shotbolt et al. 2005, 2006; Filstrup et al. 2010). In young reservoirs, the unsupported/supported ^{210}Pb boundary may not be reached. Furthermore, the 1963 ^{137}Cs peak, produced by atmospheric nuclear bomb testing, and often used for dating validation, may not be present. Additionally, eutrophication indicators such as organic matter, nutrient concentrations, and isotopes of carbon ($\delta^{13}\text{C}$) and nitrogen ($\delta^{15}\text{N}$) may be more difficult to interpret in reservoirs than in natural lakes. This is because artificial water bodies age faster and have different mixing regimes and morphological characteristics (Shotbolt et al. 2005, 2006).

There are many reservoirs in the south-central USA, which have allegedly undergone water quality changes, but long-term monitoring data are generally lacking, making it difficult to attribute cause and identify solutions. In these situations, paleolimnological methods are employed for inferring past changes (Costa-Böddeker et al. 2012). For example, Filstrup et al. (2010) utilized a suite of geochemical variables in Lake Waco, Texas to reconstruct water quality. The study successfully documented the eutrophication history of the reservoir. Organic carbon, organic nitrogen, total phosphorus and $\delta^{15}\text{N}$ all increased, whereas C:N and N:P decreased. Although the study was successful, confounding factors such as large allochthonous organic matter (OM) loads, differing OM degradation rates and changes in external N sources complicated interpretation of the sediment isotope profile from the reservoir. The authors concluded that more research was needed to understand isotopes in reservoir sediments before interpretations of past water quality can be made (Filstrup et al. 2010).

Arkansas is officially known as the “Natural State” for its clear lakes, streams and abundance of wildlife. In particular, northwest Arkansas (NWA) is dotted with caves, sinkholes, losing streams, plateaus and valleys typical of a karst-dominated terrain. The northern region is home to the Buffalo River, the first National River to be designated in the United States. It remains one of the few rivers in the lower 48 states that have not been dammed. The northern region has experienced sustained economic and population growth since the 1970s, with increasing development pressure (United States Census Bureau 2000). Because soils are thin and poorly developed, typical of karsted landscapes, agricultural production has been limited to animal production. As of 2010, Arkansas was ranked second and third in chicken and turkey production, respectively, and 24th in hogs and pigs in the USA (USDA-NASS 2010).

Approximately 55 % of Arkansas’ animal production (primarily poultry) was confined to northwest Arkansas (Slaton et al. 2004). Feeding operations from these activities generated large quantities of manure, which were land-applied (Haggard et al. 2003) and later identified as major sources of nutrient loading to water bodies, as expressed by increased concentrations of nitrogen and phosphorus in streams (Davis and Bell 1998). In addition to agriculture, the human population quadrupled between 1963 (100,000) and 2007 (400,000) in NWA, placing stresses on land and water resources. Forest cover decreased from 518 to 492 km² between 1999 and 2006, while urban land use increased from 15.5 to 46.6 km² during the same period. Total phosphorus in the water column of Beaver Reservoir, the site of this study, increased from 40 $\mu\text{g L}^{-1}$ in the 1970s to 60 $\mu\text{g L}^{-1}$ in the 2000s, while total nitrogen averaged 800 $\mu\text{g L}^{-1}$ over that time. Annual appearance of the taste-and-odor compound 2-Methyl Iso-borneol (MIB) in Beaver Reservoir during recent summers is hypothesized to be from deteriorating water quality and consequent toxic algae proliferation. Citizens, water managers and other stakeholders continue to express concern over water quality in this reservoir.

We analyzed geochemistry and diatoms in an 83-cm sediment core from Beaver Reservoir to determine if changes in water quality were preserved in the sediments and if these changes were correlated with anthropogenic activities in the watershed. We hypothesized that biological and chemical indicators of eutrophication in the sediments would increase in response to these water quality changes.

Site description

Beaver Reservoir (94°12'N, 36°18'W) is a large, multi-purpose reservoir that was built in 1963 in the Upper White River Basin of northwest Arkansas (Fig. 1). The reservoir has a surface area of 114 km² and a maximum depth of 63 m. The catchment area is 3,087 km², of which 57 % is in forest, 32 % in agriculture, and 5 % in urban use (Galloway and Green 2006). The air temperature ranges from 0 to 38 °C, with a 30-year mean annual temperature of 20 °C (NOAA 2009). The main inflows to the reservoir are the White River and War Eagle Creek (Green 1996; Haggard and Green 2002; Galloway and Green 2006). River flows generally reflect precipitation. From 1964 to 2012, total annual precipitation averaged 1,169 mm, with 30 % falling during spring. During the



Fig. 1 Map of beaver reservoir, watershed boundaries and population centers within the *watershed*. Fayetteville and Rogers are not within the watershed boundary, but drain major

streams that discharge into the reservoir. An 83-cm core was retrieved from the mid-reservoir zone indicated by the *black oval*

same period, average annual flow into the reservoir was 558 CFS with highest flows in the fall and spring. The reservoir is warm monomictic, with an average water residence time of 1.5 years (National Eutrophication Survey 1977). Average pool elevation is 341 m above sea level and average daily release (hydropower and spillway) is $3.4 \times 10^6 \text{ m}^3$. Thermal stratification usually begins in May and complete mixing begins in October or November, with continued circulation throughout the winter season.

Materials and methods

Sediment coring, dating and geochemical analysis

An 83-cm sediment core was extracted from Beaver Reservoir using a UWITEC[®] percussion gravity corer on 18 September 2007. The core was taken in the middle zone of the reservoir in approximately 25 m of water. The collected core had approximately 20 cm of water above the sediment surface, which was gently

removed using a tube and a 20-ml syringe. The core was then sectioned at 1-cm intervals from top to bottom, and samples were placed in Ziplock bags and kept on ice during transport to the lab.

The sediment chronology was determined by measuring ^{210}Pb , ^{137}Cs and ^{226}Ra in the Radioisotope Laboratory at the University of Manitoba, Canada. Sixteen sample depths were counted for ^{210}Pb and six samples were measured for ^{137}Cs . Five to 15 grams of sediment were sealed in petri dishes and “aged” for approximately 30 days to enable ^{226}Ra to equilibrate with daughters ^{214}Pb and ^{214}Bi . After aging, samples in the sealed dishes were counted on a gamma spectrometer (HyperpureGe detector) to determine ^{210}Pb , ^{137}Cs and ^{226}Ra activities. Unsupported ^{210}Pb activity in each sample was determined by subtracting ^{226}Ra activity (i.e. supported ^{210}Pb activity) from total ^{210}Pb activity at each level. Sediment ages were calculated using the constant rate of supply (CRS) model (Appleby and Oldfield 1978; Oldfield and Appleby 1984) and the constant initial concentration (CIC) model (Robbins 1978). Age errors for the CRS model were propagated using first-order approximations and calculated according to Binford (1990). Because the reservoir was only 45 years old at the time of core collection, and reservoir sediments had accumulated for far <6 half-lives of ^{210}Pb , we could not measure the total integrated, unsupported ^{210}Pb (mBq cm^{-2}) that should have accumulated at the core site. We therefore used a novel approach to estimate the total unsupported ^{210}Pb value (A_0) expected at the site, had sediment been accumulating before the date of reservoir construction. That was accomplished as shown in the equations below, using the integrated unsupported ^{210}Pb measured in the core (A_{core}), i.e. since reservoir construction, to account for the “missing” unsupported ^{210}Pb that would have accumulated before the construction of the reservoir ($A_{\text{pre-reservoir}}$)

$$A_{\text{pre-reservoir}} = A_{\text{core}} \left(e^{(\text{reservoir age} * \text{decay constant})} - 1 \right)^{-1}$$

$$A_{\text{pre-reservoir}} = 616.7 \left(e^{(44.8 * 0.03114)} - 1 \right)^{-1} = 203.3$$

$$A_0 = A_{\text{core}} + A_{\text{pre-reservoir}} = 616.7 + 203.3 = 820$$

We assessed organic matter (OM) content, carbon and nitrogen concentration and stable isotopic values in organic matter to determine past lake productivity.

OM content was determined by loss on ignition (LOI) following Heiri et al. (2001). Approximately 2 g of dry sediment was combusted at 550 °C in a muffle furnace for 4 h. The difference between the dry weight and ash weight is expressed as % LOI. Organic matter $\delta^{13}\text{C}$ was determined to identify organic matter sources. Samples were acidified with 10 % HCl to remove carbonates and dried overnight in a low-temperature oven. Nitrogen and carbon concentrations and their stable isotopes were measured on an elemental analyzer and Thermo-Finnigan Delta Plus IR-MS, respectively, at the University of Arkansas Stable Isotope Laboratory (UASIL, Fayetteville, Arkansas). Measured organic $\delta^{13}\text{C}$ were adjusted to correct for anthropogenic changes in atmospheric carbon isotopic values in the past two centuries (Suess effect), using the equation from Verburg (2007). Total P in sediment was measured using the persulfate digestion method (EPA 365.4).

Diatom analysis

Sediments were treated with 30 % H_2O_2 and 10 % HCl at 80 °C. Once the organic matter was oxidized, test tubes were refrigerated overnight at 4 °C to stop the oxidation by H_2O_2 and allow diatoms to settle. The diatom slurry was spiked with microspheres (Batterbee and Kneen 1982) and mounted on round cover slips. The cover slips were dried overnight at room temperature and then mounted on slides using Naphrax[®]. Diatom taxonomy was determined using Krammer and Lange-Bertalot (1986–1991). At least 300 diatom valves were identified per sample. Statistical analyses were performed using SYSTAT[®] 12.0, with statistical significance set at $p < 0.05$ (Sokal and Rohlf 1995). Pearson correlation was used to test significant relationships between variables.

Results

Core characteristics

The 83-cm-long sediment core consisted of brown gyttja with no observable differences in texture. Excess ^{210}Pb activity was greatest at the surface (76 Bq g^{-1}) and declined exponentially with accumulated dry mass ($r^2 = 0.62$, $p < 0.001$, Fig. 2a). The




Fig. 2 a Activity of unsupported ²¹⁰Pb (filled circles) and ¹³⁷Cs (open circles) versus accumulated dry mass and (b) depth-age relationship using the CRS (filled circles) and CIC (open circles) models from beaver reservoir sediment, Arkansas. Error bars indicate one SD

mean sedimentation rate was calculated as 0.332 g cm⁻² a⁻¹ and was obtained by dividing total accumulated dry mass (14.93 g cm⁻²) by the age of the reservoir (45 year). The CRS model estimated variable sedimentation in the core (0.2–0.8 g cm⁻² a⁻¹). The ¹³⁷Cs activity remained relatively constant throughout the core, with a small increase in activity at the base (Fig. 2a).

Sediment chronology and bulk density

Sediment ages obtained using the CRS and CIC models are presented in Table 1. Mean dates provided by both models were similar (Table 1, Fig. 2b). The 95 % confidence intervals, however, were larger for the CIC model. For example, the date of the basal layer ranged from 1946–1984 using the CIC model, compared to 1962–1964 using the CRS model. The larger confidence intervals for CIC dates suggest that the

Table 1 Comparison of dates and 95 % confidence intervals from the beaver reservoir core, using the CRS and CIC ²¹⁰Pb models

Max Depth (cm)	CRS			CIC		
	Low	Mean	High	Low	Mean	High
1	2005.2	2007.1	2009.1	2006.8	2007.1	2007.4
6	2002.9	2004.9	2006.9	2003	2004.5	2006
11	2000.6	2002.6	2004.6	1999.4	2002	2004.6
16	1998.4	2000.4	2002.5	1996.2	1999.8	2003.4
21	1996.3	1998.4	2000.5	1993.5	1997.9	2002.4
26	1993.8	1995.9	1998.1	1990.5	1995.9	2001.2
31	1990.9	1993.1	1995.4	1986.9	1993.4	1999.9
36	1988.5	1990.8	1993.1	1983.4	1991	1998.5
41	1985.7	1988.1	1990.4	1979.9	1988.5	1997.2
46	1982.3	1984.7	1987.2	1976.3	1986.1	1995.9
51	1979	1981.5	1984	1972.6	1983.5	1994.5
56	1975.6	1978.1	1980.7	1968.6	1980.8	1992.9
61	1972.1	1974.7	1977.3	1964.8	1978.1	1991.5

variable sedimentation rates estimated by the CRS model more accurately reflect sedimentation in the core. The sediment core had an average bulk density of 0.18 ± 0.04 g cm⁻³, but showed a decreasing trend from 1975 to 1998, followed by an increasing trend from 1999 to 2007 (Fig. 3).

Geochemical and biological indicators

LOI increased from 5.4 % at the bottom of the core to 7.0 % at the top (Fig. 4a). Although the range was small (1.6 %), the increase was significant (r² = 0.42, p < 0.001). The C:N molar ratio averaged 7.42 ± 0.43, but did not show a clear trend with age (Fig. 4b). Similar to LOI, diatom abundance increased significantly from the bottom to the top of the core (Fig. 4c, r² = 0.12, p = 0.003). The diatom *Aulacoseira ambigua* (Grunow) Simonsen was the most abundant taxon throughout the core and reached a maximum relative abundance of 40 % around 2004 (Fig. 4d).

Total nitrogen and total carbon were positively correlated (r = 0.80, n = 83, p < 0.001), suggesting both were bound in the organic matter. Total nitrogen displayed a general increasing trend from 1963 to 2007 (Fig. 5a, r² = 0.29, p < 0.001). The minimum TN was 2.4 mg g⁻¹, a value that occurred several




Fig. 3 Sediment bulk density of the beaver reservoir sediment profile




Fig. 4 **a** % LOI, **b** carbon to nitrogen (C:N) ratio, **c** diatom abundance and **d** *Aulacoseira ambigua* abundance versus date in the beaver reservoir sediment profile

times along the core. The maximum TN of 3.4 mg g^{-1} was measured at the top of the core in 2007. Total phosphorus concentration also increased significantly between 1963 and 2007 (Fig. 5b, $r^2 = 0.40$, $n = 24$, $p < 0.001$). TP was approximately 0.10 mg g^{-1} from the 1960s to 1990, then generally $>0.20 \text{ mg g}^{-1}$ to 2007. Maximum TP measured was 0.55 mg g^{-1} in 1998 (Fig. 5b). TN:TP decreased significantly from 1963 to 2007 (Fig. 5c, $r^2 = 0.28$, $p < 0.01$). Between 1963 and 1990 the TN:TP was generally >20 , with

values <15 after 1990 (Fig. 5c). Both TN and TP were significantly correlated to human population increase (Fig. 6, both $p < 0.001$).

Stable isotopes of carbon ($\delta^{13}\text{C}$) and nitrogen ($\delta^{15}\text{N}$)

The uncorrected and corrected $\delta^{13}\text{C}$ values were highly variable from 1963 to 1982 (Fig. 7a), with three periods of lighter $\delta^{13}\text{C}$, around 1968, 1976 and 1982. From 1982 to 2007, the $\delta^{13}\text{C}$ values are much less variable (Fig. 7a), with corrected values generally remaining between -25 and -24 ‰ (Fig. 7a). The $\delta^{15}\text{N}$ averaged 7.45 ± 0.44 ‰ (Fig. 7b). There was a general decline from 8 to 7 ‰ from ca. 1964 to 1990, but no apparent trend from 1990 to 2007 (Fig. 7b). The $\delta^{15}\text{N}$ did not display an overall trend from the bottom to the top of the core and was not correlated with any other measured variable.

Discussion

Results from our study suggest that sediment cores can serve as useful archives for trophic status change in reservoirs, particularly in areas where natural lakes do not exist. The use of cores to assess change and ascribe these changes to watershed activity is largely based on the ability to demonstrate that the sediment was not disturbed and to reliably date the sediments. Once this has been accomplished, observed changes in the core, such as increased nutrients, can be correlated to watershed activities. Several models for sediment dating with ^{210}Pb and ^{137}Cs have been developed. These are discussed extensively in Robbins (1978) and Appleby and Oldfield (1978). Here, we used our ^{210}Pb dates cautiously to place observed geochemical changes in the sediment core in a general temporal context. We did so because our results illustrate some of the challenges in using radioisotopes to date reservoir sediments.

In our study, we compared dates and sedimentation rates estimated by the CIC and CRS models. Ideally the core should extend below the unsupported/supported ^{210}Pb horizon. The core in our study did not extend below the unsupported/supported ^{210}Pb horizon because of the reservoir's young age, so we estimated the "missing" component of the total unsupported ^{210}Pb inventory from the ^{210}Pb inventory

Fig. 5 **a** Total nitrogen, **b** total phosphorus, **c** nitrogen to phosphorus (N:P) ratio versus date in the beaver reservoir sediment profile





Fig. 6 Human population in the beaver reservoir watershed versus concentrations of **(a)** total nitrogen (TN) and **(b)** total phosphorus (TP) in the beaver reservoir sediment profile

in the core and the reservoir's age. Again, under ideal circumstances, the ¹³⁷Cs profile should display a down-core increase in activity to a sharp peak, below which there is a rapid decrease to zero. This peak corresponds to the 1963 maximum in atmospheric atomic bomb testing and can sometimes be used to verify dates generated from the ²¹⁰Pb model. Whereas there was a slight increase in ¹³⁷Cs at the core base in

the Beaver Reservoir core, there was no defined peak and pre-1963 sediments were not obtained. Absence of the peak hinders the ability to independently verify the ²¹⁰Pb dates and highlights the challenge of sediment dating in young reservoirs. Despite this challenge, the exponential decline in excess ²¹⁰Pb with accumulated dry mass indicates orderly deposition of sediments since reservoir construction. Although we found general agreement between CRS and CIC dates, the larger confidence intervals for CIC dates indicate variable sedimentation rates over time and the CRS model confirmed these variable sedimentation rates. Because the CRS and CIC models yielded similar dates and indicated changing sedimentation rates over time, we have confidence that the changes we see in sediment variables over time are placed in a reasonable temporal framework.

To demonstrate that the sediments provide an accurate representation of past conditions, we correlated historical information on land use change in the watershed and changes in environmental conditions in the reservoir. Historical information on the drainage basin was sparse. The best proxies for environmental change were the growth in population, which quadrupled between 1963 (100,000) and 2007 (400,000), and the increase in annual broiler (poultry) production, from 259,850 × 10³ to 1,175,900 × 10³ over the same period (<http://nass.usda.gov>). Our results and those of others (Davis and Bell 1998) suggest that these anthropogenic activities impacted nutrient conditions and biological productivity in the reservoir. For example, the increase in sediment TN and TP was significantly correlated to population growth. Biological activity also increases up-core, as indicated by the significant increases in LOI, diatom abundance and higher percentages of *A. ambigua*, an indicator of eutrophic conditions (Interlandi et al. 1999; Zalat and




Fig. 7 a Uncorrected $\delta^{13}\text{C}$ (solid line) and $\delta^{13}\text{C}$ corrected for the Suess effect (dotted line) and (b) $\delta^{15}\text{N}$ versus date in the beaver reservoir sediment profile

Vildary 2007). Between 1993 and 1995, a study was conducted to examine the amount of poultry waste production, and potential impacts on water quality (Haggard et al. 2003). The study revealed that the Beaver Reservoir basin produced about 200,000 metric tons a^{-1} of poultry litter, with most being land-applied as fertilizer to meet forage N requirements (Haggard et al. 2003). Land-applied poultry litter had been cited for increasing N and P in northwest Arkansas (NWA) streams (Petersen 1992; Haggard et al. 2003). Observed increases in sediment TN and TP may have been a consequence of animal manure application and subsequent stream runoff to the reservoir.

The impact of increased TP was evident in the TN:TP ratio, which fell from an average of 27 between 1963 and 1990 to 11 between 1990 and 2007 (Fig. 5). Smith (1982) reported that TN:TP mass ratio < 15 indicates N limitation, whereas $\text{N:P} > 16$ indicates P limitation. Nitrogen limitation often leads to cyanobacteria proliferation. It appears that the reservoir switched from P limitation between 1963 and 1990 to N limitation from 1990 to 2007. Over the last 10 years, the reservoir has experienced episodic outbreaks of taste and odor compounds that have been linked to cyanobacterial proliferation in other reservoirs (Watson 2003; Youngstead 2005). In addition to poultry litter application, other aspects of human population

growth probably contributed to increased nutrient concentrations. In NWA, unpaved roads were much more common in the 1960s and 1970s compared to today. The decrease in sediment bulk density in the mid-1980s might reflect reduced allochthonous sediment from unpaved roads. Regional water authorities are reducing sediment input from the watershed by paving streets. Our study suggests that the recent change in sediment type was driven by an increase in the input of nutrients, not inorganic material.

Algal OM and terrestrial OM (C_3 plants) have overlapping $\delta^{13}\text{C}$, which range from -25 to -30 ‰. Thus, $\delta^{13}\text{C}$ is often used in combination with the C:N ratio to differentiate organic matter source. Typically, algal-derived organic matter has a C:N atomic ratio from 2 to 10, whereas terrestrially derived organic matter has C:N from 20 to 80 (Meyers and Lallier-Vergès 1999). In our study, the C:N atomic ratio was consistently < 10 throughout the core and the $\delta^{13}\text{C}$ was in the range expected for algal-derived OM, with $\delta^{13}\text{C}$ values between -23.5 and -30 ‰. The study suggests that algal organic matter was the predominant source of organic matter to the sediments, providing further evidence for deteriorating water quality and enhanced biologic production in the reservoir.

The $\delta^{13}\text{C}$ can also be used to infer past productivity because of the carbon isotope fractionation that occurs during photosynthesis. Periods of very high productivity

generally lead to greater incorporation of ^{13}C into the OM as algae draw down the supply of ^{12}C . Greater $\delta^{13}\text{C}$ values are thus found in the sediment core organic matter (Hodell and Schelske 1998; Mizutani and Wada 1982). The trend toward higher organic matter $\delta^{13}\text{C}$ during periods of greater productivity was not observed at Beaver Reservoir. Several factors could account for the lack of an increasing trend in organic matter $\delta^{13}\text{C}$ accompanying enhanced productivity (Gu et al. 2004, 2006; Rosenmeier et al. 2004; Woodward et al. 2012). Atmospheric CO_2 equilibrates with aqueous CO_2 , thereby bringing down the carbon isotopic value in water and algal OM, obscuring inferences about productivity (Verburg 2007). Sewage effluent may also influence the $\delta^{13}\text{C}$ of sediments. Rosenmeier et al. (2004) attributed depleted ^{13}C in recent sediments of Lake Petén Itza, Guatemala, to sewage effluent, because sewage OM is low in ^{13}C relative to OM produced by phytoplankton. Given that our Beaver Reservoir core was retrieved downstream from a wastewater treatment plant, it is possible that the carbon isotope values in the sediment OM reflect, in part, sewage effluent input.

Photosynthesis by lacustrine algae results in a stable carbon isotope fractionation of about 20 ‰ (Hecky and Hesslein 1995). Fractionation, however, is reduced and $\delta^{13}\text{C}$ of OM is consequently greater when productivity is high and $\text{CO}_{2\text{aq}}$ is in short supply. Respiration on the other hand contributes CO_2 with low ^{13}C content to the water. Because most lakes with a significant supply of allochthonous OM are net heterotrophic (respiration exceeds photosynthesis), and supersaturated with CO_2 (Cole et al. 1994), there is perhaps an abundant supply of $^{12}\text{CO}_2$ for photosynthesis. This supply may mask the effects of high productivity that would be observed in lakes where $^{12}\text{CO}_2$ is in short supply.

Increased $\delta^{15}\text{N}$ has also been used as an indicator of algal productivity in numerous studies (Altabet and Francois 1994; Teranes and Bernasconi 2000; Filstrup et al. 2009). As epilimnetic dissolved inorganic nitrogen (DIN) pools become progressively enriched with ^{15}N from preferential assimilation and hypolimnetic transport of ^{14}N , phytoplankton discriminate less against ^{15}N (Hodell and Schelske 1998). The $\delta^{15}\text{N}$ in our study did not demonstrate a trend of enrichment as observed in natural lakes experiencing increased primary productivity. Rather, the $\delta^{15}\text{N}$ in our core decreased, but only from 8 to 7 ‰ between 1963 and

1990. The lack of a large change in $\delta^{15}\text{N}$, in response to greater productivity in the reservoir, may have been a consequence of several factors. Only 26 % of the variation in $\delta^{15}\text{N}$ ($r^2 = 0.26$, $n = 37$) was accounted for by TN concentration, suggesting other factors influenced both N concentration and its isotopic ratio. Because N was non-limiting before 1990, phytoplankton discriminated against ^{15}N and readily incorporated ^{14}N into their biomass, thus displaying relatively low $\delta^{15}\text{N}$ up to 1990. When the reservoir became N-limited after 1990, the $\delta^{15}\text{N}$ of sediments fluctuated, perhaps in response to availability and source of nitrogen. For example, in the late 1990s and 2000, $\delta^{15}\text{N}$ was as high as 8.5 ‰, which suggests lower N concentrations and thus greater incorporation of ^{15}N by phytoplankton (Fig. 3). Conversely, low $\delta^{15}\text{N}$ ca. 2002 and 2003 suggests greater overall N concentrations and more discrimination against ^{15}N . The $\delta^{15}\text{N}$ of 8.5 ‰ may also be a reflection of sewage effluent because sewage $\delta^{15}\text{N}$ ranges from 10 to 20 ‰.

The shift in the N:P ratio towards N limitation may have led to greater cyanobacterial proliferation in the reservoir which, in turn, could have affected N dynamics in two ways. First, because cyanobacteria fix N directly from the atmosphere, the amount of ^{14}N in the water column available for incorporation into algal biomass could also increase. Although the phenomenon has not been shown in freshwater systems, approximately 52 % of fixed nitrogen can become available for algal uptake in oceans (Mullholland et al. 2006). Second, cyanobacteria $\delta^{15}\text{N}$ values are generally close to zero, reflective of their substrate, air. Therefore, high abundances of cyanobacteria will pull $\delta^{15}\text{N}$ values down toward zero. This may explain the lower $\delta^{15}\text{N}$ ca. 2002 and 2003. Lower $\delta^{15}\text{N}$, possibly initiated by cyanobacterial fixation, is further supported by the observation that 2002 had the highest level of the cyanobacterially-produced taste and odor compound 2-Methyl-Isoborneol on record. Other mechanisms, perhaps N mineralization (Bernasconi et al. 1997), may have led to lower $\delta^{15}\text{N}$ in sediments, whereas selective loss of N through denitrification or immobilization (Talbot and Laerdal 2000; Teranes and Bernasconi 2000; Savage et al. 2004) led to higher $\delta^{15}\text{N}$. Grantz et al. (2012) demonstrated that denitrification accounts for significant losses of N in reservoirs. These losses further suggest a significant impact on the $\delta^{15}\text{N}$ signature of sediments.

Conclusions

We demonstrated that sediments of Beaver Reservoir recorded increasing concentrations of N and P through time and chronicled a shift from P limitation to N limitation around 1990. These increases coincided with increases in human population in the watershed, land application of poultry manure and documented increases in N and P concentrations in streams. The study also documented a significant increase in organic matter concentration derived from algal sources, based on the C:N ratio and $\delta^{13}\text{C}$ values. Stable isotope analysis of C and N suggested several processes influenced the $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ signature of the sediments. Further studies are needed to better understand the behavior of $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ in reservoir sediments to more effectively enable their use in historical water quality investigations. Despite these challenges, sediment studies could provide valuable information on historical water quality in reservoirs.

Acknowledgments We thank the Beaver Water District for assistance with fieldwork, and Anna Nottemeir. We also thank Troy Munhofen and Lanayah Turley, NSF-funded REU students who assisted with stable isotope measurements during the summer of 2009. Funding for this project was provided by the Arkansas Water Resource Center, Beaver Water District and a USGS 104B Grant.

References

- Altabet MA, Francois R (1994) Sedimentary nitrogen isotopic ratio as a recorder for surface ocean nitrate utilization. *Global Biogeochem Cycles* 8:103–116
- Appleby PG, Oldfield F (1978) The calculation of ^{210}Pb dates assuming a constant rate of supply of unsupported ^{210}Pb to the sediment. *Catena* 5:1–8
- Batterbee RW, Kneen MJ (1982) The use of electronically counted microspheres in absolute diatom analysis. *Limnol Oceanogr* 27:184–188
- Bernasconi SM, Barbieri A, Simona M (1997) Carbon and nitrogen isotope variations in sedimenting organic matter in Lake Lugano. *Limnol Oceanogr* 42:1755–1765
- Cole J, Caraco NF, Kling GW, Kratz TK (1994) Carbon dioxide supersaturation in the surface waters of lakes. *Science* 265:1568–1570
- Costa-Böddeker S, Bennion H, de Jesus TA, Albuquerque ALS, Figueira RCL (2012) Paleolimnologically inferred eutrophication of a shallow tropical, urban reservoir in Southeast Brazil. *J Paleolimnol* 48:751–766
- Davis JV, Bell RW (1998) Water quality assessment of the ozark plateaus study unit, Arkansas, Kansas, Missouri, and Oklahoma: nutrients, bacteria, organic carbon, and suspended sediment in surface water, 1993–95. US Geological Survey Water Resources Investigations Report Publication 98–4164
- Filstrup CT, Scott JT, Lind OT (2009) Allochthonous organic matter supplements and sediment transport in a polymictic reservoir determined using elemental and isotopic ratios. *Biogeochemistry* 96:87–100
- Filstrup CT, Scott JT, White DJ, Lind OT (2010) Use of element and isotopic compositions to record the eutrophication of a polymictic reservoir in central Texas, USA. *Lake Reservoir Manag* 15:25–39
- Flynn WW (1968) The determination of low levels of Po-210 in environmental materials. *Anal Chim Acta* 43:221–227
- Galloway JM, Green WR (2006) Analysis of ambient conditions and simulation of hydrodynamics and water quality characteristics in Beaver Lake, Arkansas, 2001 through 2003. US Geological Survey Scientific Investigations Report 2006–5003
- Grant EM, Kogo A, Scott JT (2012) Partitioning whole-lake denitrification using in situ dinitrogen gas accumulation and intact sediment core experiments. *Limnol Oceanogr* 57:925–935
- Green WR (1996) Eutrophication trends inferred from hypolimnetic dissolved oxygen dynamics within selected white river reservoirs, northern Arkansas—southern Missouri 1974–1994. US Geological Survey Scientific Investigations Report 1996–4096
- Gu B, Schelske CL, Hodell DA (2004) Extreme ^{13}C enrichments in a shallow hypereutrophic lake: implications for carbon cycling. *Limnol Oceanogr* 49:1152–1159
- Gu B, Chapman AD, Schelske CL (2006) Factors controlling seasonal variations in stable isotope composition of particulate organic matter in a soft water eutrophic lake. *Limnol Oceanogr* 51:2837–2848
- Haggard BE, Green WR (2002) Simulation of hydrodynamics, temperature and dissolved oxygen in Beaver Lake, Arkansas, 1994–1995. US Geological Survey Scientific Investigations Report 2002–4116
- Haggard BE, Moore PA, Chaubey I, Stanley EH (2003) Nitrogen and phosphorus concentrations and export from an Ozark Plateau catchment in the United States. *Biosyst Eng* 86:75–85
- Hecky RE, Hesslein RH (1995) Contributions of benthic algae to lake food webs as revealed by stable isotope analysis. *JN Am Benthol Soc* 14:631–653
- Heiri O, Lotter AF, Lemcke G (2001) Loss on ignition as a method for estimating organic and carbonate content in sediments: reproducibility and comparability of results. *J Paleolimnol* 25:101–110
- Hodell DA, Schelske CL (1998) Production, sedimentation, and isotopic composition of organic matter in Lake Ontario. *Limnol Oceanogr* 43:200–214
- Interlandi SJ, Kilham SS, Theriot EC (1999) Response of phytoplankton to varied resource availability in large lakes of the Greater Yellowstone Ecosystem. *Limnol Oceanogr* 44:668–682
- Meyers PA, Lallier-Vergès E (1999) Lacustrine sedimentary organic matter records of Late Quaternary paleoclimates. *J Paleolimnol* 21:345–372

- Mizutani M, Wada H (1982) Effect of high atmospheric CO₂ concentrations on $\delta^{13}\text{C}$ of algae. *Origins Life Evol B* 12:377–390
- Mullholland MR, Bernhardt PW, Heil CA, Bronk DA, O'Neil JM (2006) Nitrogen fixation and release of fixed nitrogen by *Trichodesmium* spp. in the Gulf of Mexico. *Limnol Oceanogr* 51:1762–1776
- National Eutrophication Survey (1977). Report on beaver, table rock, and bull shoals reservoirs, arkansas and taneycomo reservoir, missouri. Working Paper No. 480. USEPA, regions VI and VII
- NOAA National Oceanic Atmospheric Association (2009) Accessed June 2009. Monthly Statistics Available at <http://www.srh.noaa.gov/tsa/climate/normrain.html>
- Oldfield F, Appleby PG (1984) Empirical testing of ²¹⁰Pb-dating models for lake sediments. In: Haworth EY, Lund JWG (eds) *Lake sediments and environmental history*. University of Minnesota, Minneapolis, pp 93–124
- Petersen JC (1992) Trends in stream water quality data in Arkansas during several time periods between 1975 and 1989. US Geological Survey Water-Resources Investigation Report, Publication 92–4044
- Révész K, Haiping Qi (2006) Determination of the δ (¹⁵N/¹⁴N) and δ (¹³C/¹²C) of total N and C in solids: RSIL lab code 1832, chap. C5 of Révész K and Coplen TB (eds) *Methods of the reston stable isotope laboratory: Reston Virginia, US Geological Survey, Techniques and Methods, book 10, sec. C, chap. 5, 31*
- Robbins JA (1978) Geochemical and geophysical applications of radioactive lead isotopes. In: Nriago JP (ed) *Biogeochemistry of lead*. North Holland, Amsterdam, pp 285–393
- Rosenmeier MF, Brenner M, Kenney W, Whitmore TJ, Taylor CM (2004) Recent eutrophication in the southern basin of Lake Petén Itzá, Guatemala: human impact on a large tropical lake. *J Paleolimnol* 511:161–172
- Savage C, Leavitt PR, Elmgren R (2004) Distribution and retention of effluent nitrogen in surface sediments of a coastal bay. *Limnol Oceanogr* 49:1503–1511
- Shotbolt LA, Thomas AD, Hutchinson SM (2005) The use of reservoir sediments as environmental archives of catchment inputs and atmospheric pollution. *Prog Phys Geogr* 29:337–361
- Shotbolt L, Hutchinson SM, Thomas AD (2006) Sediment stratigraphy and heavy metal fluxes to reservoirs in the southern Pennine uplands, UK. *J Paleolimnol* 35:305–322
- Slaton AN, Brye RK, Daniels MB, Daniel CT, Norman JR, Miller MD (2004) Nutrient input and removal trends for agricultural soils in nine geographic regions in Arkansas. *J Environ Qual* 33:1606–1615
- Smith VH (1982) The nitrogen and phosphorus dependence of algal biomass in lakes. An empirical and theoretical analysis. *Limnol Oceanogr* 27:1101–1112
- Sokal RR, Rohlf FJ (1995) *Biometry: the principles and practice of statistics in biological research*, 3rd edn. W. H. Freeman and Company, NY
- Talbot MR, Laerdal T (2000) The late Pleistocene-Holocene paleolimnology of Lake Victoria, East Africa, based upon elemental and isotopic analyses of sedimentary organic matter. *J Paleolimnol* 23:141–164
- Teranes JL, Bernasconi SM (2000) The record of nitrate utilization and productivity limitation provided by d15 N values in lake organic matter—a study of sediment trap and core sediments from Baldeggersee, Switzerland. *Limnol Oceanogr* 45:801–813
- Torres IC, Inglett PW, Brenner M, Kenney WF, Reddy RK (2012) Stable isotope ($\delta^{13}\text{C}$ and $\delta^{15}\text{N}$) signatures of sediment organic matter in subtropical lakes of different trophic status. *J Paleolimnol* 147:693–706. doi:10.1007/s10933-012-9593-6
- United States Census Bureau (2000) Population by county and place census 2000: Available at <http://www.Census.gov/population/www/census2000/respop.html>
- USDA-NASS (2010) 2010 Census of agriculture: state and county profiles. Available at http://www.nass.usda.gov/Statistics_by_State/Arkansas/index.asp
- Verburg P (2007) The need to correct for the Suess effect in the application of $\delta^{13}\text{C}$ in sediment of autotrophic Lake Tanganyika, as a productivity proxy in the Anthropocene. *J Paleolimnol* 37:591–602
- Watson SB (2003) Cyanobacterial and eukaryotic algal odor compounds: signals or byproducts? A review of their biological activity. *Phycologia* 42:332–350
- Woodward AC, Polito PA, Beilman WD (2012) Carbon and nitrogen stable isotope ratios in surface sediments from lakes of western Ireland: implications for inferring past lake productivity and nitrogen loading. *J Paleolimnol* 47:167–184
- Youngstead NW (2005) Factors that influence phosphorus, filamentous cyanobacteria and odor in McDaniel Lake, a Southwest Missouri water supply reservoir, 1983–2002. *Lake Reserv Manag* 21:453–464
- Zalat A, Vildary SS (2007) Environmental change in Northern Egyptian Delta lakes during the late Holocene, based on diatom analysis. *J Paleolimnol* 37:273–299